



Diagnosis of pseudoxanthoma elasticum in a patient with discrete skin lesions[☆]

Dear Editor,

Pseudoxanthoma elasticum (PXE) is an autosomal recessive genetic metabolic condition characterized by aberrant calcification with the fragmentation of elastic fibers in the dermis, retina and tunica intima of arteries.¹ We present the case of a patient with ocular alterations and slight cutaneous signs as a debut form of PXE, which was confirmed by skin biopsy.

A 33-year-old woman with a history of systemic lupus erythematosus was referred because an ocular fundus examination revealed angiod streaks associated with decreased foveolar brightness and diffuse pigmentary changes called "peau d'orange" (Fig. 1). Physical examination revealed two 2 mm whitish papules on the lateral cervical fold (Fig. 2). The patient was normotensive with peripheral and symmetrical pulses present. Calcium, phosphorus, and lipid profile values were normal. Histological examination of the cervical lesions revealed thickened and fragmented dermal elastic fibers. Von Kossa staining was positive for calcium deposition in elastic fibers (Fig. 3), confirming the diagnosis of PXE.

The prevalence of PXE varies between 1/25,000 and 1/100,000, with female predominance. It is caused by a mutation in the ABCC6 gene, located on chromosome

16, which encodes a transmembrane transport protein MRP6.² The pathophysiology involves reduced levels of the anti-mineralization factor inorganic pyrophosphate (PPi), dysfunctional extracellular calcium homeostasis, and ectopic mineralization of tissues rich in elastic fibers. There is a considerable phenotypic heterogeneity that could explain the late diagnosis of the clinical case presented.³

Although cutaneous findings usually represent the first clinical sign, they are typically subtle and not evident until the second or third decade.³ The first cutaneous sign is small, discrete, yellowish papules at the lateral side of the neck, axillae, antecubital and popliteal fossae. These papules coalesce to form plaques of corrugated and inelastic appearance.^{2,3} Despite the innocuous appearance of these skin changes, they may reveal significant ocular and vascular involvement. Subsequently, it is associated with redundant skin in flexural folds, generally in the axillae and groin.⁴ In some patients with angiod streaks, a biopsy of healthy axillary or scar skin may reveal histological findings characteristic of PXE.⁴ Electron microscopy of the skin reveals bulky mineral deposits that disrupt and break elastic fibers in the mid-dermis.²

The first visible changes on funduscopy are pigment irregularities with a "peau d'orange" appearance that typically precedes angiod streaks by one to eight years. Angiod streaks originate from the optic disc and radiate outwards as brownish-grey irregular lines, histopathological findings show breaks of the calcified and thickened Bruch's membrane. The latter predisposes to secondary choroidal




Figure 1 Fundus photograph of the right eye. Note the "peau d'orange" appearance of the fundus temporal to the macula (arrow) and barely visible angiod streaks radiate outwards from the optic disc giving the appearance of a "spider web" (arrowhead). Fluorescein angiography of the left eye demonstrated the angiod streaks with hyperfluorescent edges radiating from the optic nerve up to the midperiphery (arrow).

[☆] Study conducted at the Hospital Barros Luco Trudeau, Santiago, Chile.




Figure 2 (A) Left cervicolateral skin area with two barely perceptible whitish papules, 2 mm in diameter, marked with a blue pencil (arrows). (B) Dermoscopy of one of the whitish papules marked with blue pencil.




Figure 3 (A) Epidermis without alterations, dermis with increased and fragmented elastic fibers, which are thickened and with increased basophilia (Hematoxylin & eosin, 40×). (B) Von Kossa stain positive for calcium deposition in elastic fibers (Von Kossa, 40×).

neovascularization with the consequent risk of hemorrhage and blindness in the later stages of the disease.⁵ Although highly suggestive, angioid streaks have also been reported in patients with hemoglobinopathies, Paget's disease, and Ehler-Danlos syndrome.²

Manifestations of vascular involvement include loss of peripheral pulses, claudication, hypertension, myocardial infarction and ischaemic or hemorrhagic strokes. These findings reflect the mineralization of the middle and intimal layers of small and medium-caliber arteries.^{2,3}

We present this case given the unusual clinical presentation and highlight the importance of cutaneous alterations in the diagnostic confirmation of this entity.

Financial support

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions

Catalina Jahr: Approval of the final version of the manuscript; composition of the manuscript; collection, analysis, and interpretation of data; participation in the design of the study; critical review of the literature; critical review of the manuscript.

Valentina Vera: Approval of the final version of the manuscript; collection, analysis, and interpretation of data; critical review of the manuscript.

Roberto Bustos: Approval of the final version of the manuscript; collection, analysis, and interpretation of data; critical review of the manuscript.

José Contreras: Approval of the final version of the manuscript; collection, analysis, and interpretation of data; critical review of the manuscript.

Conflicts of interest

None declared.

References

- Gliem M, Zaeytijd JD, Finger RP, Holz FG, Leroy BP, Issa PC. An update on the ocular phenotype in patients with pseudoxanthoma elasticum. *Front Genet.* 2013;4:14.
- Germain DP. Pseudoxanthoma elasticum. *Orphanet J Rare Dis.* 2017;12:85.
- Uitto J, Jiang Q, Váradí A, Bercovitch LG, Terry SF. Pseudoxanthoma elasticum: diagnostic features, classification and treatment options. *Expert Opinion on Orphan Drugs.* 2014;2:567–77.
- Brown SJ, Talks SJ, Needham SJ, Taylor AEM. Pseudoxanthoma elasticum: biopsy of clinically normal skin in the investigation of patients with angiod streaks. *Br J Dermatol.* 2007;157:748–51.
- Georgalas I, Tservakis I, Papaconstantinou D, Kardara M, Koutsandrea C, Ladas I. Pseudoxanthoma elasticum, ocular manifestations, complications and treatment. *Clinical and Experimental Optometry.* 2011;94:169–80.

Catalina Jahr  ^{a,*}, Valentina Vera  ^a, Roberto Bustos  ^b, José Contreras 

^a Department of Dermatology, Hospital Clínico de la Universidad de Chile, Santiago, Chile

^b Department of Dermatology, Hospital Barros Luco Trudeau, Santiago, Chile

* Corresponding author.

E-mail: catalina.jahr@gmail.com, catalina.jahr@ug.uchile.cl (C. Jahr).

Received 27 July 2021; accepted 30 July 2021;

Available online 20 April 2023

<https://doi.org/10.1016/j.abd.2021.07.012>

0365-0596/ © 2023 Published by Elsevier España, S.L.U. on behalf of Sociedade Brasileira de Dermatologia. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Generalized nodular tinea profunda in an immunosuppressed patient caused by *Trichophyton rubrum*☆



Dear Editor,

We report the case of a 65-year-old man with generalized subcutaneous nodules with intense pruritus for 1 year (Fig. 1). He had a history of diabetes mellitus for 7 years, hypertension and chronic kidney disease for 5 years, and Bullous Pemphigoid (BP) for 2 years. Systemic glucocorticoid (oral prednisone 25 mg twice a day) was prescribed to treat his BP for more than 1 year.

After admission, blood test for fungal glucan was 213 pg/mL (the normal is lower than 60 pg/mL), which indicated a deep fungal infection. Blood culture for fungi was negative, and no lymphadenopathy was detected by ultrasonography. The light microscopy showed epidermal hyperplasia, dermal abscess, and infiltration of neutrophils, lymphocytes, epithelioid cells, and scattered multinucleated giant cells (Fig. 2A). Intracellular hyphae were observed in multinucleated cells in the granuloma (Fig. 2B). The skin sample was also sent for Next Generation Sequencing (NGS) to identify the pathogen. The NGS reported *Trichophyton rubrum* nucleotide sequences (Cover rate: 0.0199%) in DNA extracted from the skin specimen. Considering the patient's clinical and histological manifestations, and the notable high sequencing reads compared to a negative control, we established the diagnosis of generalized nodular tinea profunda caused by *Trichophyton rubrum*. Oral therapy with 250 mg terbinafine per day was initiated and the nodules regressed completely after 3 months.

Trichophyton rubrum often causes superficial dermatomycosis, such as tinea manus, tinea pedis and tinea corporis. But in very few cases, *T. rubrum* penetrate into the dermis

☆ Study conducted at the Department of Dermatology, The Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, China.