

Figure 2 Orthohyperkeratosis, absence of granular cell layer and mild acanthosis are observed in the epidermis. In the upper dermis is a mild perivascular lymphocytic infiltrate (Hematoxylin & eosin, $\times 200$).

intellectual content; Collection, analysis, and interpretation of data; Critical review of the literature; Approval of the final version of the manuscript.

Conflicts of interest

None declared.

References

- Mishra K, Jandial A, Gupta K, Prakash G, Malhotra P. Ichthyosis: a harbinger of lymphoma. *BMJ Case Rep.* 2018;2018, bcr2018224229.
- Word AP, Cayce R, Pandya AG. Beware of underlying malignancy: acquired ichthyosis. *Am J Med.* 2014;127:202-4.

İrfan Yavaşoğlu ^a, Atakan Turgutkaya ^{a,*}, Canten Tataroğlu ^b, Ali Zahit Bolaman ^a

^a *Division of Hematology, Aydin Adnan Menderes University Medical Faculty, Aydin, Turkey*

^b *Division of Pathology, Aydin Adnan Menderes University Medical Faculty, Aydin, Turkey*

* Corresponding author.

E-mail: atakanturgutkaya@yahoo.com.tr (A. Turgutkaya).

Received 19 June 2023; accepted 16 August 2023

Available online 14 July 2024

<https://doi.org/10.1016/j.abd.2023.08.020>

0365-0596/ © 2024 Sociedade Brasileira de Dermatologia.

Published by Elsevier España, S.L.U. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Confocal reflectance microscopy in basal cell carcinoma associated with nevus sebaceous: case report[☆]



Dear Editor,

Nevus sebaceous is a congenital benign hamartoma of the skin. Its most common complication is transformation into other tumors, usually benign.^{1,2} However, due to the potential for malignancy, early diagnosis and treatment are essential.^{3,4}

While dermoscopy allows the analysis of the epidermis to the mid-dermis, reflectance confocal microscopy (RCM) uses an 830-nm diode laser as a monochromatic and coherent light source which penetrates, between 200 and 300 μm , providing images at the cellular level that resemble histopathology, offering detailed morphological analysis of the different skin layers up to the papillary dermis.⁵

There are few reports in the literature regarding typical findings of nevus sebaceous in RCM.⁶⁻⁸ Articles describing lesions associated with basal cell carcinoma are even less frequent.^{9,10} The authors present a scenario of common

dermoscopic characteristics in nevus sebaceous associated with basal cell carcinoma, highlighting changes in confocal reflectance microscopy of the nevus sebaceous, scarcely described in the literature to date.

A 41-year-old male patient with no personal or family history of skin cancer, was treated for a lesion present since childhood on the right forehead with changes in texture and slow growth over the years.

Clinically there was a pearly-yellow plaque, with unclear borders, on an erythematous base and telangiectasias on the periphery, and yellowish papules in its upper region. Palpation showed a slightly verrucous texture (Fig. 1A).

Dermoscopy showed, in the lower region, round and oval, whitish and whitish-yellow uniformly aggregated structures in a cobblestones pattern, with telangiectasias on the periphery. There were arboriform vessels in the central region, which are typically associated with basal cell carcinoma. The upper region showed a group of rounded whitish-yellow papules with central umbilication and crown vessels (Fig. 1B).

RCM, carried out with VivaScope[®] 1500 (Lucid Inc. Rochester, NY, USA) showed, in the dermis, typical findings of basal cell carcinoma: tumor islands with peritumoral clefting, dark silhouettes and, on the periphery, palisaded cells and dilated tortuous vessels (Fig. 2). At the dermal-epidermal junction and papillary dermis, central tube-shaped structures stood out, with sebaceous gland

[☆] Study conducted at the Hospital A.C. Camargo Cancer Center, São Paulo, SP, Brazil.




Figure 1 (A) Macroscopy of a pearly-yellow plaque with unclear borders on an erythematous base (*) with telangiectasias on the periphery (1) and yellowish papules (2) on the upper region. (B) Dermoscopy ($\times 10$ magnification) showing round whitish-yellow structures in a cobblestone pattern (3), telangiectasias (4), arboriform vessels (5) and rounded whitish-yellow papules with central umbilication and crown vessels (6).




Figure 2 Confocal reflectance microscopy of the papillary dermis showing basal cell carcinoma with individual images measuring $0.5 \times 0.5 \text{ mm}^2$ and a mosaic measuring $8 \times 8 \text{ mm}^2$. (A and B) Tumor island (*) with peritumoral clefting (1) and dark silhouettes (2).

lobes in the surrounding area, filled with aggregates similar to fish ova, typical of nevus sebaceous (Fig. 3).

Two incisional biopsies were performed with a 4 mm punch. Histopathology revealed in the lower region, superficial basal cell carcinoma and, in the central region, superficial and nodular basal cell carcinoma, both associated with nevus sebaceous (Fig. 4).

The patient underwent surgery with margin control and primary closure. He has been followed in the Cutaneous Oncology service for six months, with no signs of recurrence.

Nevus sebaceous is a congenital benign hamartoma of the skin consisting of numerous malformed sebaceous glands, degenerated hair follicles and ectopic apocrine glands, most often located on the face and scalp.¹ Two-thirds of the lesions are present from birth and one-third develop in early childhood.² Its most common complication is transformation into other tumors, more frequently benign ones, firstly trichoblastoma, followed by papillary syringocystadenoma.³ Among malignant tumors, the most common is basal cell carcinoma, which develops in less than 1% of cases.⁴ Due to the




Figure 3 Confocal reflectance microscopy of the papillary dermis of the nevus sebaceous with individual images measuring $0.5 \times 0.5 \text{ mm}^2$ and mosaics measuring $8 \times 8 \text{ mm}^2$. (A and B) Central tube-shaped structures (*), with sebaceous gland lobes in the surrounding area (1), filled with aggregates similar to fish ova (2).




Figure 4 Histopathology. (A) Nodular basal cell carcinoma with an adenoid pattern, in association with a nevus sebaceous. (B) Superficial and nodular basal cell carcinoma in association with a nevus sebaceous (Hematoxylin & eosin, A $\times 100$ and B $\times 200$).

potential for malignant transformation, early diagnosis and treatment are essential.

While dermoscopy allows the analysis of the epidermis up to the superficial reticular dermis, reflectance confocal microscopy (RCM) uses an 830-nm diode laser as a monochromatic and coherent light source. The penetration depth, between 200 and 300 μm , provides images at the cellular level that resemble histopathology, offering a detailed morphological analysis of the different skin layers up to the papillary dermis.⁵

There are few reports in the literature to date regarding the typical findings of nevus sebaceous in RCM. Descriptions approximate those of sebaceous hyperplasia in RCM.⁶⁻⁸ A single study identified changes in nevus sebaceous on confocal imaging in different age groups.⁹ The article showed that, under the age of ten, hypoplastic sebaceous glands and juvenile hair follicles can be seen in these lesions. From ten to 59 years, the sebaceous glands at the dermal-epidermal junction resemble bunches of grapes and, in the superficial dermis, structures similar to tubes or loops can be seen in the center, which correspond to the dilation of the seba-

ceous duct,⁷ with sebaceous gland lobes resembling fish ova in the surrounding area and verrucous or papillomatous hyperplasia in the dermis. Above the age of 60, papillomatous hyperplasia predominates on RCM examination.¹⁻³

RCM reports describing the characteristics of nevus sebaceous associated with those of basal cell carcinoma, are even less frequent at present, with a single case report of papillary syringocystadenoma and basal cell carcinoma arising from a previous nevus sebaceous.¹⁰

Therefore, new studies are necessary so that more typical structures of nevus sebaceous associated with basal cell carcinoma are described on RCM.

Financial support

None declared.

Authors' contributions

Ingrid Priscila Ribeiro Paes Ferraz: Approval of the final version of the manuscript; drafting and editing of the manuscript; review of the literature; critical review of the manuscript.

Gustavo Carvalho: Approval of the final version of the manuscript; drafting and editing of the manuscript; review of the literature; critical review of the manuscript.

Juliana Casagrande Tavoloni Braga: Approval of the final version of the manuscript; drafting and editing of the manuscript; review of the literature; critical review of the manuscript.

Rafaela Brito de Paula: Approval of the final version of the manuscript; drafting and editing of the manuscript; review of the literature; critical review of the manuscript.

André Molina: Approval of the final version of the manuscript; drafting and editing of the manuscript; review of the literature; critical review of the manuscript.






Conflicts of interest

None declared.

References

1. Lima TFM, Melo MN, Baeninger LG, Neto AG, Simião AL. Nevus sebaceous with malignant transformation treated with Mohs surgery: a case report. *Surg Cosmet Dermatol*. 2020;12:204-7.

2. Constant E, Davis DG. The premalignant nature of the sebaceous nevus of Jadassohn. *Plast Reconstr Surg*. 1972;50:257-9.
3. Idriss MH, Elston DM. Secondary neoplasms associated with nevus sebaceous of Jadassohn: a study of 707 cases. *J Am Acad Dermatol*. 2014;70:332-7.
4. Cribier B, Scrivener Y, Grosshans E. Tumors arising in nevus sebaceous: a study of 596 cases. *J Am Acad Dermatol*. 2000;42:263-8.
5. Rezza GG, Casagrande JT. Microscopia Confocal (MC). In: Rezza GG, Casagrande JT, editors. *Atlas de Microscopia Confocal na Dermatologia*. São Paulo: Lemar; 2016. p. 21-3.
6. Rao L, Lin EY, Wang WJ, Huang XW. Detection of sebaceous gland hyperplasia with dermoscopy and reflectance confocal microscopy. Preprint. 2022:1-10.
7. Propperova I, Langley RG. Reflectance-mode confocal microscopy for the diagnosis of sebaceous hyperplasia in vivo. *Arch Dermatol*. 2007;143:134.
8. González S, White WM, Rajadhyaksha M, Anderson RR, González E. Confocal imaging of sebaceous gland hyperplasia in vivo to assess efficacy and mechanism of pulsed dye laser treatment. *Lasers Surg Med*. 1999;25:8-12.
9. Jiang Q, Chen H, Ma Ling, Huang M, Xia Y, Chen L. Characteristic analysis of sebaceous nevus using dermoscopy and reflectance confocal microscopy. *Chinese Journal of Dermatology*. 2018:523-5.
10. Jiang J, Chen Y, He Q, Yang J, Zhang Z, Yang H, et al. Syringocystadenoma papilliferum and basal cell carcinoma arising in nevus sebaceous. *Clin Cosmet Investig Dermatol*. 2022;15:2021-6.

Ingrid Priscila Ribeiro Paes Ferraz  ^{a,*},
Gustavo Carvalho  ^b,
Juliana Casagrande Tavoloni Braga  ^b,
Rafaela Brito de Paula  ^b, André Molina  ^b

^a *Emergency Department, Hospital A.C. Camargo Cancer Center, São Paulo, SP, Brazil*

^b *Department of Cutaneous Oncology, Hospital A.C. Camargo Cancer Center, São Paulo, SP, Brazil*

* Corresponding author.

E-mail: ingrid.ferraz@accamargo.org.br (I.P. Ferraz).

Received 23 April 2023; accepted 25 September 2023
Available online 23 August 2024

<https://doi.org/10.1016/j.abd.2023.09.011>

0365-0596/ © 2024 Sociedade Brasileira de Dermatologia.

Published by Elsevier España, S.L.U. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).